

Guideline



CCHMC Trauma Service Guidelines

Title: CRAFFT Screening

Effective Date: 06/2018

Number: TR-25

Page: 1 of 4

1.0 SCOPE

- 1.1. Care of the Trauma Services Patient at CCHMC.

2.0 DEFINITIONS

- 2.1. **CRAFFT**: A validated tool for pre-adolescent / adolescent substance abuse screening for children age 12 to 17 years. (CRAFFT: **C**ar, **R**elax, **A**lone, **F**orget, **F**riends, **T**rouble).

3.0 GUIDELINE

- 3.1. CRAFFT screen should be completed for all injured patients age 12 to 17 years.
 - 3.1.1. Substance abuse screening is required by the American College of Surgeons (ACS) for all trauma centers.
 - 3.1.2. Complete CRAFFT tool as per instructions (see form below).
 - 3.1.2.1. Access CRAFFT doc-flow sheet in EPIC to complete screening and document score.
 - 3.1.2.2. A paper copy of CRAFFT screen is also available and may be given to the patient to be completed.
 - 3.1.2.3. If paper screen is used this data must then be entered into the patient's medical record.
- 3.2. A Social Service consult is indicated for any patient who scores 1 – 6 for brief intervention.
 - 3.2.1. Nursing to enter Social Service request in EPIC if indicated.
 - 3.2.2. Social Service will complete brief intervention and document recommendations in patient medical record.

The CRAFFT Screening Interview

Begin: “I’m going to ask you a few questions that I ask all my patients. Please be honest. I will keep your answers confidential.”

Part A

During the PAST 12 MONTHS, did you: **No** **Yes**

- | | | |
|--|--------------------------|--------------------------|
| 1. Drink any <u>alcohol</u> (more than a few sips)?
(Do not count a few sips of alcohol taken during family or religious events) | <input type="checkbox"/> | <input type="checkbox"/> |
| 2. Smoke any <u>marijuana</u> or <u>hashish</u> ? | <input type="checkbox"/> | <input type="checkbox"/> |
| 3. Use <u>anything else</u> to get high? (“ <u>anything else</u> ” includes other illegal drugs, prescription or non-prescription medicines or things that people sniff or “huff”) | <input type="checkbox"/> | <input type="checkbox"/> |

For clinic use only: Did the patient answer “yes” to any questions in Part A?

No



Ask CAR question only, then stop

Yes



Ask all 6 CRAFFT questions

Part B

- | | No | Yes |
|---|--------------------------|--------------------------|
| 1. Have you ever ridden in a CAR driven by someone (including yourself) who was “high” or had been using alcohol or drugs? | <input type="checkbox"/> | <input type="checkbox"/> |
| 2. Do you ever use alcohol or drugs to RELAX , feel better about yourself, or fit in? | <input type="checkbox"/> | <input type="checkbox"/> |
| 3. Do you ever use alcohol or drugs while you are by yourself, or ALONE ? | <input type="checkbox"/> | <input type="checkbox"/> |
| 4. Do you ever FORGET things you did while using alcohol or drugs? | <input type="checkbox"/> | <input type="checkbox"/> |
| 5. Do your FAMILY or FRIENDS ever tell you that you should cut down on your drinking or drug use? | <input type="checkbox"/> | <input type="checkbox"/> |
| 6. Have you ever gotten into TROUBLE while you were using alcohol or drugs? | <input type="checkbox"/> | <input type="checkbox"/> |


*Two or more **YES** answers on the CRAFFT suggest a serious problem and need for further assessment.

CONFIDENTIALITY NOTICE:

The information recorded on this page may be protected by special federal confidentiality rules (42 CFR Part 2), which prohibit disclosure of this information unless authorized by specific written

© John R. Knight, MD, Boston Children’s Hospital, 2014.
All rights reserved. Reproduced with permission.
For more information, contact ceasar@childrens.harvard.edu

1. Show your patient his/her score on the graph



2. Talking Points for Clinician Counseling

No Use

- Not using drugs and alcohol is a smart decision for your health. You should be proud of yourself. If that ever changes, I hope you trust me enough that we can talk about it.

Any Use

- My recommendation as your doctor is not to use drugs or alcohol *at all*.

CRAFFT score ≥ 2

- I am quite concerned about you. Let's plan a follow-up meeting in about a month.

3. Counseling Points for All

- Alcohol and drugs can affect brain development, which continues into your mid-20s.
- Early alcohol and drug use greatly increases your risk for developing addiction, major depression, anxiety disorders, and psychotic thinking.
- Alcohol can hurt and scar your liver; marijuana can hurt your lungs
- Alcohol and drugs can cause car crashes. Marijuana impairs driving just as much as alcohol does, and its effects last longer.

4. Give each patient the Contact for Life

<http://www.sadd.org/contract.htm#cf1>

The Center for Adolescent Substance Abuse Research,
(CeASAR) at Boston Children's Hospital
(617) 355-5433 www.ceasar.org
For more information, contact ceasar@childrens.harvard.edu

4.0 REFERENCES

- 4.1 CDC. Youth Risk Behavior Surveillance- United States,. (2009). MMWR Recomm Rep, 59.
- 4.2 NIAAA Alcohol screening and brief intervention for youth: A practitioner's guide. (2011). Accessed May 1, 2015, at www.niaaa.nih.gov/YouthGuide).
- 4.3 SAMHSA. Results from the 2010 National Survey on Drug Use and Health: Summary of National Findings. (2011). Rockville, MD: SAMHSA.

5.0 APPROVALS

All revisions of this guideline are approved by the Trauma Service Department. This guideline is reviewed every three years or sooner if deemed necessary. Policy authority for this document resides with the Trauma Service Department. This guideline is approved by the Trauma Service Manager and the Director of Trauma Services.

HISTORY	
Original Date	
	09/2009
Revision Date	
	05/15
Review Date	
	06/18