

Procedure

CCHMC Trauma Service Procedure	<i>Procedure Number</i>	TR-03
Criteria for Trauma Team Activation	<i>Effective Date</i>	9/2017
	<i>Page</i>	1 of 4

1.0 SCOPE

- 1.1. Trauma Resuscitation Team is activated based upon information from the field, a referring hospital or on the patient's condition on presentation to the Emergency Department (ED).
- 1.2. If at any time there is question regarding status of the patient, the Trauma Resuscitation Team should be activated.
- 1.3. A three tiered response system allows for appropriate utilization of personnel.
 - 1.3.1. Trauma Stat for the unstable and most seriously injured child.
 - 1.3.2. Trauma Alert for the stable but seriously injured child.
 - 1.3.3. Trauma Evaluation (Eval) for the child with potential for injury and needs rapid evaluation.
- 1.4. Notification for Trauma Stat and Trauma Alert occurs via hospital paging system. All members are simultaneously notified of an incoming trauma patient.

2.0 DEFINITIONS

- 2.1. **Trauma Stat:** highest activation which mobilizes a 22 member team from various disciplines within the hospital in order to manage the unstable and most seriously injured child.
- 2.2. **Trauma Alert:** mobilization of team except for anesthesia, OR personnel and radiology who are not required to attend; used to manage the stable but seriously injured child.
- 2.3. **Trauma Eval:** lowest level of activation, initiated with the arrival of a stable, previously evaluated child or when a child arrives with a trauma mechanisms which is minor in nature.
- 2.4. **TBSA:** Total Body Surface Area
- 2.5. **Normal vital signs for age:**

Age	Pulse Beats/min	Respirations Breaths/min	Avg. Systolic BP	Avg. Diastolic BP
Premature	120 – 170	40 – 70	55 – 75	35 – 45
0 – 3 months	100 – 150	35 – 55	65 – 85	45 – 55
3 – 6 months	90 – 120	30 – 45	70 – 90	50 – 65
6 – 12 months	90 – 120	25 – 40	80 – 100	55 – 65
1 – 3 years	70 – 110	20 – 30	90 – 105	55 – 70
3 – 6 years	65 – 110	20 – 25	95 – 110	60 – 75
6 – 12 years	60 – 95	14 – 22	100 – 120	60 – 75
12+ years	55 – 85	12 – 18	110 – 135	65 – 85

Behman, RE, Kliegman, RM & Jenson, HB. (2003). *Nelson Textbook of Pediatrics*. Saunders

3.0 PROCEDURES

- 3.1. Activation Criteria for **Trauma Stat:**
 - 3.1.1. Any penetrating wound of the head, neck or trunk
 - 3.1.1.1. Gunshot wound (GSW) to the head, neck or trunk or extremities proximal to the elbow/knee
 - 3.1.2. Tachycardia and / or poor perfusion or unexplained tachycardia (no significant pain or crying as a source)
 - 3.1.3. Blood given prior to the patient's arrival
 - 3.1.4. Hypotension
 - 3.1.5. 40 mL/kg bolus prior to arrival
 - 3.1.6. Respiratory difficulty as evidenced by
 - 3.1.6.1. Significant increase or decrease in respiratory rate
 - 3.1.6.2. Significant retractions or grunting
 - 3.1.6.3. Patient intubated prior to arrival
 - 3.1.6.4. Unable to maintain or difficult airway
 - 3.1.7. Glasgow Coma Score (GCS) \leq 8
 - 3.1.8. GCS deterioration by 2
- 3.2. Activation Criteria for **Trauma Alert:**
 - 3.2.1. Evidence of abdominal injury

Procedure

CCHMC Trauma Service Procedure	<i>Procedure Number</i>	TR-03
Criteria for Trauma Team Activation	<i>Effective Date</i>	9/2017
	<i>Page</i>	2 of 4

- 3.2.1.1. Without hemodynamic compromise
- 3.2.1.2. Distended and / or tender abdomen
- 3.2.1.3. Abdominal bruising or seat belt mark

3.2.2. GCS 9 – 13

3.2.3. Spinal cord injury with neurologic deficit

3.2.4. Two or more proximal long bone fractures

3.2.5. Burns > 15% TBSA

3.2.6. Ejection from vehicle

3.2.7. Significant vascular injury including amputation of limb proximal to wrist or ankle

3.2.8. Emergency Department discretion

3.3 Activation Criteria for **Trauma Eval:**

3.3.1. Motor vehicle collision

3.3.2. Struck or run over by motor vehicle (pedestrian or bike)

3.3.3. Fall greater than 10 feet

3.3.4. Any mechanism deemed to place the patient at risk for multi-system injury

3.3.5. Any patient immobilized with a backboard and/or cervical collar

3.3.6. Partial or full thickness burns between 5% and 14% TBSA

3.3.7. Any burn less than 5% TBSA requiring immediate pain management

3.3.8. Any GSW (non-BB) distal to the knee or elbow

3.3.9. Application of a tourniquet prior to arrival and hemodynamically stable

3.4. **Criteria for patients seen at an outside facility prior to arrival at CCHMC**

3.4.1. The trauma team should be activated for any patient who meets criteria regardless of prior evaluation at an outside facility.

3.5 Responding Personnel to Trauma Resuscitations

Personnel	Stat	Alert	Eval
Dr. Right (surgeon)	X	X	
Nurse Right (ED)	X	X	
Dr Left (pediatric resident)	X	X	X
Nurse Left (Trauma Core RN)	X	X	
Anesthesiologist	X		
Respiratory Therapist	X	X	
Attending Surgeon	X		
Physician Team Leader (ED)	X	X	X
PICU Fellow	X		
Nurse Team Leader (Trauma Core RN)	X	X	X
Medication Nurse (PICU and ED)	X	X	
Nurse Liaison (MPS)	X	X	
Fluid (paramedic)	X	X	X
OR Nurse	X		
PCA	X	X	X
Security	X	X	
X-ray Tech	X	X	
Chaplains	X	X	
Social Services	X		
Radiologist	X		
Child Life Specialist	X	X	X
Service Center	X	X	

4.0 LIST OF ATTACHED FORMS

4.1. N/A

5.0 REFERENCES

Procedure

CCHMC Trauma Service Procedure	<i>Procedure Number</i>	TR-03
Criteria for Trauma Team Activation	<i>Effective Date</i>	9/2017
	<i>Page</i>	3 of 4

5.1. Resources for Optimal Care of the Injured Patient: 2006; American College of Surgeons Committee on Trauma; Chicago, Illinois.

6.0 APPROVALS

All revisions of this procedure are approved by the Trauma Services Department. This procedure is reviewed every three years or sooner if deemed necessary. Authority for this document resides with the Trauma Services Department. This procedure is approved by the Trauma Service Manager and the Director of Trauma Services.

HISTORY	
Original Date	
05/95	
Revision Date	
7/00, 9/06, 10/10, 1/13, 3/14, 2/15, 10/16, 10/17	
Review Date	
10/16	

Procedure

CCHMC Trauma Service Procedure	<i>Procedure Number</i>	TR-03
Criteria for Trauma Team Activation	<i>Effective Date</i>	9/2017
	<i>Page</i>	4 of 4

Trauma Stat:

- Any penetrating injury to head, neck or trunk
- GSW to the extremities proximal to the elbow / knee
- Respiratory difficulty as evidenced by one or more of the following:
 - Significant increase or decrease in respiratory rate
 - Significant retractions or grunting
 - Patient intubated prior to arrival
 - Unable to maintain or difficult airway
- Tachycardia and / or poor perfusion or unexplained tachycardia (no significant pain or crying as a source)
- Hypotension
- Blood given prior to the patient's arrival
- 40 mL/kg bolus given prior to arrival
- Glasgow Coma Score (GCS) ≤ 8
- GCS deterioration by 2

Normal Vital Signs Table

Age	Pulse Beats/min	Respirations Breaths/min	Avg. Systolic BP
Premature	120 – 170	40 – 70	55 – 75
0 – 3 months	100 – 150	35 – 55	65 – 85
3 – 6 months	90 – 120	30 – 45	70 – 90
6 – 12 months	90 – 120	25 – 40	80 – 100
1 – 3 years	70 – 110	20 – 30	90 – 105
3 – 6 years	65 – 110	20 – 25	95 – 110
6 – 12 years	60 – 95	14 – 22	100 – 120
12+ years	55 – 85	12 – 18	110 – 135

Behman, RE, Kliegman, RM & Jenson, HB. (2003). *Nelson Textbook of Pediatrics*. Saunders

OR Resuscitation:

- Full arrest with pre-hospital signs of life following a non-cranial, penetrating chest or abdominal injury
- Penetrating injury unresponsive to 40 mL/kg fluid administration
- At the discretion of the ED and surgical attending

Trauma Alert:

- Evidence of abdominal injury
- Without hemodynamic compromise
- Distended and / or tender abdomen
- Abdominal bruising or seatbelt mark
- GCS 9 – 13
- Spinal cord injury with neurologic deficit
- Two or more proximal long bone fractures
- Ejection from automobile
- Partial or full thickness burn of $>15\%$ TBSA
- Significant vascular injury including amputation of limb proximal to wrist or ankle
- Emergency Department discretion

Trauma Evaluation

- Motor vehicle collision
- Struck or run over by motor vehicle (pedestrian or bike)
- Fall greater than 10 feet
- Any mechanism deemed to place the patient at risk for multi-system injury
- Any patient immobilized with a backboard and/or cervical collar
- Partial or full thickness burn between 5% and 14% TBSA
- Any burn less than 5% requiring immediate pain management
- Any GSW (non-BB) distal to the knee or elbow
- Tourniquet application prior to arrival and hemodynamically stable

*****The trauma team should be activated for any patient who meets criteria regardless of prior evaluation at an outside facility*****